

Use Ballpoint Pen Only

"Authorization is hereby given to dispense the Generic or therapeutic equivalent unless otherwise indicated by the words -- NO SUBSTITUTE"

Do Not Use Abbreviations	Q.D., QD, q.d., qd (Daily) MS or MSO4 (Morphine Sulfate)	U or u (Unit) IU (international units)	Q.O.D., QOD, q.o.d., qod (every other day) MgSO4 (Magnesium Sulfate)
Spine Preoperative Orders Items with evidence indicated by ^			
At Orthopedist's Office	☐ Patient Weight _____ Patient Height _____ ☐ Assess for specialty bed if weight greater than 500 lbs ☐ Give prescription for Chlorhexidine wash and instruct patient to use wash on day prior to surgery AND morning of surgery		
Attending Surgeon:	Physician assistant:		
Patient Status	☐ Inpatient on arrival to SDS (Prior Authorization, IP Only CPT, or stay anticipated longer than 2 MN) ☐ Outpatient		
Enter Surgical Procedure	SPINE:		
Vital Signs	☒ Vital signs per protocol		
Nursing Orders	☒ If MRSA Screening is positive, place patient on CONTACT PRECAUTIONS ☒ Hair removal to be performed by practitioner ☒ Type & Screen unless patient has on an arm band ☐ Type & Cross 2 units Packed Red Blood Cells ☒ Cleanse operative site with Chlorhexidine wipes^ ☐ Apply Knee length intermittent Pneumatic Compression Device to Bilateral Lower Extremities ☐ Capillary Blood Glucose on arrival to SDS ☐ Stat PT/INR ☐ Teach preoperative incentive spirometry 10 x per hour		
IV Fluids	☐ IV Lactated Ringers at _____ ml/hr ☐ IV _____ at _____ ml/hr		
Medications	For patients on home beta blockers, their normal oral dose should be taken within the 24 hours prior to surgery ☐ Home Beta Blocker is currently contraindicated by _____ Prophylactic Antibiotics:^ ☐ Cefazolin (Ancef®) 2 gram IV (3 gram IV for patients 120 kg or above) on arrival to OR. Redose every 4 hours during surgery. Cefazolin contraindicated with allergy to Cephalosporins or severe Penicillin allergy (anaphylaxis, urticaria, breathing problems, or angioedema)—Please order Clindamycin instead. If patient has allergy to clindamycin as well as cephalosporin allergy or severe penicillin allergy please use vancomycin ☐ Clindamycin (Cleocin®) 900 mg IV over 30 minutes x 1 dose. Redose every 6 hours during surgery. ☐ Vancomycin weight-based dose (see below)- instead of cefazolin or clindamycin ☒ Nozin application upon arrival to SDS ☒ If nares PCR results are positive for MRSA, then give Vancomycin weight-based dose (see below) just prior to the start of either Cefazolin or Clindamycin as ordered above^ in addition to cefazolin or clindamycin Vancomycin weight-based dosing: <i>< 80 kg: 1g IV infused over 1 hour; 80 to 119 kg: 1.5g IV infused over 1.5 hours; 120 kg or above: 2g IV infused over 2 hours</i>		
Operating Room:	☒ Activate Knee Length Intermittent Pneumatic Compression Devices on Bilateral Lower Extremities ☐ Insert Indwelling Urinary Catheter under anesthesia after prophylactic antibiotics have been started		
PACU	☒ Vital signs per protocol in PACU ☒ Bilateral Lower Extremity IPC's ☐ Hemoglobin and Hematocrit in PACU ☐ AP/Lateral of: ☐ C-spine ☐ L-Spine		
Print Practitioner Name			
Practitioner Signature:		Date:	Time:
Drug Allergies: ☐ NKDA ☐ Other:			



169600012

SPINEPREOP



Revised 0826

Spine Preoperative Orders

Patient Full Name: _____

Patient DOB: _____ M or F

Date of Procedure: _____